arXiv:1601.01595v1 [math.CO] 7 Jan 2016

COMPOSITIONS COLORED BY SIMPLICIAL POLYTOPIC NUMBERS

DANIEL BIRMAJER, JUAN B. GIL, AND MICHAEL D. WEINER

ABSTRACT. For a given integer $d \geq 1$, we consider $\binom{n+d-1}{d}$-color compositions of a positive integer ν for which each part of size n admits $\binom{n+d-1}{d}$ colors. We give explicit formulas for the enumeration of such compositions, generalizing existing results for n-color compositions (case $d = 1$) and $\binom{n+1}{2}$-color compositions (case $d = 2$). In addition, we give bijections from the set of $\binom{n+d-1}{d}$-color compositions of ν to the set of compositions of $(d+1)\nu - 1$ having only parts of size 1 and $d+1$, the set of compositions of $(d+1)\nu$ having only parts of size congruent to 1 modulo $d+1$, and the set of compositions of $(d+1)\nu + d$ having no parts of size less than $d+1$. Our results rely on basic properties of partial Bell polynomials and on a suitable adaptation of known bijections for n-color compositions.

1. INTRODUCTION

A composition of a positive integer ν is an ordered k-tuple $(j_1, \ldots, j_k)$, for $k \geq 1$, of positive integers called parts such that $j_1 + \cdots + j_k = \nu$. We call k the number of parts.

Given a sequence of nonnegative integers $w = (w_n)_{n\in\mathbb{N}}$, we define a w-color composition of ν to be a composition of ν such that part n can take on w_n colors. If $w_n = 0$, it means that we do not use the integer n in the composition. Such colored (weighted) compositions have been considered by many authors, starting as early as 1960 (maybe even earlier), and they continue to be of current research interest, see e.g. [1, 5, 6].

If we let W_n be the number of w-color compositions of n, Moser and Whitney [8] observed that the corresponding generating functions $w(t) = \sum_{n=1}^{\infty} w_n t^n$ and $W(t) = \sum_{n=1}^{\infty} W_n t^n$ satisfy the relation

$$W(t) = \frac{w(t)}{1 - w(t)}, \text{ or equivalently, } 1 + W(t) = \frac{1}{1 - w(t)}.$$

In other words, the sequence $(W_n)_{n\in\mathbb{N}}$ is the INVERT transform of $(w_n)_{n\in\mathbb{N}}$, see [2].

A refinement of this formula, considering summations over all weighted compositions of n with exactly k parts, was given by Hoggatt and Lind [7]. They showed that the number of weighted compositions of n with exactly k parts is given by

$$c_{n,k}(w) = \sum_{\pi_k(n)} \frac{k!}{k_1! \cdots k_n!} w_1^{k_1} \cdots w_n^{k_n},$$

where the sum runs over all k-part partitions of n, that is, over all solutions of

$$k_1 + 2k_2 + \cdots + nk_n = n \text{ such that } k_1 + \cdots + k_n = k \text{ and } k_j \in \mathbb{N}_0 \text{ for all } j.$$

Note that $c_{n,k}(w)$ is precisely the partial (exponential) Bell polynomial $B_{n,k}(1!w_1, 2!w_2, \ldots)$ multiplied by a factor $k!/n!$. Given the fact that the invert transform may be written in

terms of partial Bell polynomials (via Faà di Bruno's formula), the aforementioned results can be formulated as follows.

Theorem 1. *Let $w = (w_n)_{n\in\mathbb{N}}$ be a sequence in $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. Then its invert transform*

$$W_n = \sum_{k=1}^{n} \frac{k!}{n!} B_{n,k}(1!w_1, 2!w_2, \dots) \quad \text{for } n \geq 1,$$

counts the number of w-color compositions of n. Moreover, $\frac{k!}{n!}B_{n,k}(1!w_1, 2!w_2, \dots)$ gives the number of such compositions made with exactly k parts.

This way of looking at colored compositions was recently discussed in [1] with a slightly different notation since they used the ordinary Bell polynomials instead of the exponential Bell polynomials considered here. In op. cit., the authors used this viewpoint to revisit some known examples of colored and restricted compositions.

In this paper, we study families of integer compositions colored by the *simplicial polytopic numbers* given by the sequences

$$p(d) = \left\{ \tbinom{n+d-1}{d};\ n \in \mathbb{N} \right\} \text{ for } d \in \mathbb{N}.$$

In Section 2, we make use of Theorem 1 to obtain explicit formulas for the enumeration of $p(d)$-color compositions of n, denoted by $P_n(d)$, and for the sets of restricted compositions:

$$\begin{aligned}
\mathscr{C}_{1,m}(n) &= \{\text{compositions of } n \text{ having only parts of size 1 and } m\},\\
\mathscr{C}_{\equiv 1(m)}(n) &= \{\text{compositions of } n \text{ having only parts of size congruent to 1 modulo } m\},\\
\mathscr{C}_{\geq m}(n) &= \{\text{compositions of } n \text{ having no parts of size less than } m\},
\end{aligned}$$

where m is an arbitrary integer greater than 1. In particular, we derive the identity

$$P_\nu(d) = \left|\mathscr{C}_{1,d+1}\big((d+1)\nu - 1\big)\right| = \left|\mathscr{C}_{\equiv 1(d+1)}\big((d+1)\nu\big)\right| = \left|\mathscr{C}_{\geq d+1}\big((d+1)\nu + d\big)\right|, \quad (1.1)$$

which implies that there is a one-to-one correspondence between the sets involved.

Finally, in Section 3, we provide combinatorial proofs of the identity (1.1) by suitably modifying some of the bijections given by Shapcott [9].

The results presented in this note provide a natural generalization of what is known for the set of n-color compositions (case $d = 1$) and its bijections to the set of 1-2 compositions (denoted here by $\mathscr{C}_{1,2}$), the set of odd compositions ($\mathscr{C}_{\equiv 1(2)}$), and the set of 1-free compositions ($\mathscr{C}_{\geq 2}$).

2. Enumeration formulas

We start by enumerating compositions colored by the simplicial polytopic numbers.

Proposition 2. *For $d \in \mathbb{N}$ let $p(d) = \big(p_n(d)\big)_{n\in\mathbb{N}}$ be the sequence of simplicial d-polytopic numbers $p_n(d) = \binom{n+d-1}{d}$. Then the number of $p(d)$-color compositions of n is given by*

$$P_n(d) = \sum_{k=1}^{n} \binom{n+dk-1}{n-k},$$

and $\binom{n+dk-1}{n-k}$ gives the number of such compositions having exactly k parts.

Proof. By Theorem 1, we just need to verify the identity

$$B_{n,k}(1!p_1(d), 2!p_2(d), \dots) = \frac{n!}{k!}\binom{n+dk-1}{n-k},$$

which we will prove by induction on k. For $k = 1$ and all n we have

$$B_{n,k}(1!p_1(d), 2!p_2(d), \dots) = \frac{n!}{1!}\binom{n+d-1}{d} = \frac{n!}{1!}\binom{n+d-1}{n-1}.$$

For $k > 1$, we use [3, Eq. (3k), Section 3.3] and the inductive step to get

$$\begin{aligned}
B_{n,k}(1!p_1(d), 2!p_2(d), \dots) &= \frac{1}{k}\sum_{j=0}^{n}\binom{n}{j}j!\binom{j+d-1}{d}B_{n-j,k-1}(1!p_1(d), 2!p_2(d), \dots) \\
&= \frac{1}{k}\sum_{j=1}^{n}\binom{n}{j}j!\binom{j+d-1}{d}\frac{(n-j)!}{(k-1)!}\binom{n-j+d(k-1)-1}{n-j-k+1} \\
&= \frac{n!}{k!}\sum_{j=1}^{n-k+1}\binom{j+d-1}{j-1}\binom{n-j+d(k-1)-1}{n-j-k+1} \\
&= \frac{n!}{k!}\sum_{j=0}^{n-k}\binom{j+d}{j}\binom{n-j+d(k-1)-2}{n-k-j} \\
&= \frac{n!}{k!}(-1)^{n-k}\sum_{j=0}^{n-k}\binom{-d-1}{j}\binom{-d(k-1)-k+1}{n-k-j} \\
&= (-1)^{n-k}\frac{n!}{k!}\binom{-dk-k}{n-k} = \frac{n!}{k!}\binom{n+dk-1}{n-k},
\end{aligned}$$

as claimed at the beginning of the proof. □

As discussed in the introduction, in addition to the $p(d)$-color compositions, we are also interested in the sets of restricted compositions $\mathscr{C}_{1,m}(n)$, $\mathscr{C}_{\equiv 1(m)}(n)$, and $\mathscr{C}_{\geq m}(n)$.

Proposition 3. *Let $m > 1$ be an integer. Then*

$$\left|\mathscr{C}_{1,m}(n)\right| = \sum_{j=0}^{\lfloor \frac{n}{m} \rfloor}\binom{n-(m-1)j}{j} \quad \textit{for } n \geq m.$$

Proof. Using Theorem 1 with the sequence (w_n) defined by $w_1 = w_m = 1$ and $w_j = 0$ for $j \neq 1, m$, we get $|\mathscr{C}_{1,m}(n)| = \sum_{k=1}^{n}\frac{k!}{n!}B_{n,k}(1!, 0, \dots, m!, 0, \dots)$. Now, by means of basic

properties of the partial Bell polynomials (cf. [3, Eqs. (3n) & (3n'), Section 3.3]), we get

$$\begin{aligned} B_{n,k}(1!,0,\dots,m!,0\dots) &= \sum_{\kappa=0}^{k}\sum_{\nu=\kappa}^{n}\binom{n}{\nu}B_{\nu,\kappa}(0,\dots,m!,0,\dots)B_{n-\nu,k-\kappa}(1,0,\dots) \\ &= \sum_{\kappa=0}^{k}\binom{n}{n-k+\kappa}B_{n-k+\kappa,\kappa}(0,\dots,m!,0,\dots) \\ &= \binom{n}{n-k+\frac{n-k}{m-1}}\frac{(n-k+\frac{n-k}{m-1})!}{(\frac{n-k}{m-1})!} \text{ for } n-k\equiv 0 \pmod{(m-1)}, \end{aligned}$$

and therefore

$$\left|\mathscr{C}_{1,m}(n)\right| = \sum_{\substack{1\le k\le n\\ k\equiv n \pmod{(m-1)}}} \frac{k!}{n!}\binom{n}{n-k+\frac{n-k}{m-1}}\frac{(n-k+\frac{n-k}{m-1})!}{(\frac{n-k}{m-1})!}.$$

Writing $n=q(m-1)+r$ and $k=j(m-1)+r$ with $0\le r<m-1$, we lastly obtain

$$\begin{aligned} \left|\mathscr{C}_{1,m}(n)\right| &= \sum_{j=0}^{q}\frac{(j(m-1)+r)!}{(q(m-1)+r)!}\binom{q(m-1)+r}{(q-j)m}\frac{((q-j)m)!}{(q-j)!} \\ &= \sum_{j=0}^{q}\binom{j(m-1)+r}{q-j} = \sum_{j=0}^{q}\binom{(q-j)(m-1)+r}{j} = \sum_{j=0}^{\lfloor\frac{n}{m}\rfloor}\binom{n-(m-1)j}{j}. \end{aligned}$$

□

Proposition 4. *Let $m>1$ be an integer. Then*

$$\left|\mathscr{C}_{\equiv 1(m)}(n)\right| = \sum_{j=0}^{\lfloor\frac{n}{m}\rfloor}\binom{n-(m-1)j-1}{j} \text{ for } n\ge m.$$

Proof. Consider the sequence (w_n) defined by

$$w_n = \begin{cases} 1, & \text{if } n\equiv 1 \pmod{m};\\ 0, & \text{otherwise.}\end{cases}$$

Then, for $n\ge 1$, [3, Eq. (3ℓ), Section 3.3] gives

$$\begin{aligned} B_{n,k}(1!w_1,2!w_2,\dots) &= B_{n,k}(1!,0,\dots,(m+1)!,0,\dots,(2m+1)!,0,\dots) \\ &= \sum_{\ell=0}^{k}\frac{n!}{(n-k)!\ell!}B_{n-k,k-\ell}(0,\dots,m!,0,\dots,(2m)!,0,\dots) \\ &= \sum_{\ell=0}^{k}\frac{n!}{(\frac{n-k}{m})!\ell!}B_{\frac{n-k}{m},k-\ell}(1!,2!,\dots) \text{ for } n-k\equiv 0 \pmod{m}. \end{aligned}$$

Writing $n = qm + r$ and $k = jm + r$ with $0 \le r < m$, we obtain

$$\begin{aligned}\frac{k!}{n!}B_{n,k}(1!w_1, 2!w_2, \dots) &= \sum_{\ell=0}^{k} \frac{k!}{(q-j)!\ell!} B_{q-j,k-\ell}(1!, 2!, \dots) \\ &= \sum_{\ell=0}^{k-1} \frac{k!}{(q-j)!\ell!} \frac{(q-j)!}{(k-\ell)!} \binom{q-j-1}{k-\ell-1} \quad (\text{for } j < q) \\ &= \sum_{\ell=0}^{k-1} \binom{k}{\ell} \binom{q-j-1}{k-\ell-1} = \binom{k+q-j-1}{k-1}.\end{aligned}$$

Since $\binom{k+q-j-1}{k-1} = \binom{k+q-j-1}{q-j}$, $n = qm + r$ and $k = jm + r$, we get

$$\begin{aligned}\left|\mathscr{C}_{\equiv 1(m)}(n)\right| &= \sum_{j=0}^{q} \binom{jm + r + q - j - 1}{q-j} \\ &= \sum_{j=0}^{q} \binom{(q-j)m + r + j - 1}{j} = \sum_{j=0}^{\lfloor \frac{n}{m} \rfloor} \binom{n - (m-1)j - 1}{j}.\end{aligned}$$

□

Proposition 5. *Let $m > 1$ be an integer. Then*

$$\left|\mathscr{C}_{\ge m}(n)\right| = \sum_{k=1}^{\lfloor \frac{n-1}{m-1} \rfloor} \binom{n - (m-1)k - 1}{k-1} \text{ for } n \ge m.$$

Proof. Applying Theorem 1 with the sequence (w_n) defined by $w_j = 0$ for $1 \le j < m$ and $w_j = 1$ for $j \ge m$, we get

$$\left|\mathscr{C}_{\ge m}(n)\right| = \sum_{k=1}^{n} \frac{k!}{n!} B_{n,k}(0, \dots, 0, m!, (m+1)!, \dots).$$

Now, using [3, Eqs. (3ℓ") & (3h), Section 3.3], we arrive at

$$\begin{aligned}B_{n,k}(0, \dots, m!, (m+1)!, \dots) &= \frac{n!}{(n-(m-1)k)!} B_{n-(m-1)k,k}(1!, 2! \dots) \\ &= \frac{n!}{k!} \binom{n - (m-1)k - 1}{k-1} \text{ for } (m-1)k < n.\end{aligned}$$

This completes the proof since $B_{n,k}(0, \dots, m!, (m+1)!, \dots) = 0$ for $(m-1)k \ge n$. □

As a consequence of propositions 2, 3, 4, and 5, we get the following result.

Theorem 6. *For every $d, \nu \in \mathbb{N}$, we have*

$$P_\nu(d) = \left|\mathscr{C}_{1,d+1}\big((d+1)\nu - 1\big)\right| = \left|\mathscr{C}_{\equiv 1(d+1)}\big((d+1)\nu\big)\right| = \left|\mathscr{C}_{\ge d+1}\big((d+1)\nu + d\big)\right|.$$

In other words, the set of $p(d)$-color compositions of ν is in one-to-one correspondence with the set of compositions of $(d+1)\nu - 1$ having only parts of size 1 and $d+1$, the set of

compositions of $(d+1)\nu$ *having only parts of size congruent to* 1 *modulo* $d+1$*, and the set of compositions of* $(d+1)\nu + d$ *having no parts of size less than* $d+1$*.*

3. Combinatorial bijections

Based on bijections given by Shapcott [9] for n-color compositions, in this section, we will provide bijective maps between the set of $p(d)$-color compositions of ν and the sets $\mathscr{C}_{1,d+1}\big((d+1)\nu-1\big)$, $\mathscr{C}_{\equiv 1(d+1)}\big((d+1)\nu\big)$, and $\mathscr{C}_{\geq d+1}\big((d+1)\nu+d\big)$.

For this purpose, we fix $d \in \mathbb{N}$ and consider the sets

$$A_k(\nu) = \{p(d)\text{-color compositions of } \nu \text{ with } k \text{ parts}\},$$
$$B_k(\nu) = \{\text{binary strings of length } \nu + dk - 1 \text{ with exactly } (d+1)k-1 \text{ ones}\}.$$

Proposition 7. *For any fixed* d*, there is a bijective map* $T : A_k(\nu) \to B_k(\nu)$*.*

Before we prove this proposition, we need the following lemma.

Lemma 8. *Let* $d \in \mathbb{N}$ *be fixed. For* $n \in \mathbb{N}$*,* $n \geq d$*, there is a bijection* ϕ_d *from* $\big\{1, 2, \ldots, \binom{n}{d}\big\}$ *to the set of binary words of length* n *having exactly* d *ones.*

Proof. Let $m \in \mathbb{N}$ be such that $m \leq \binom{n}{d}$. Using the fact that $\binom{n}{d} = 1 + \sum_{j=1}^{d} \binom{n-j}{d-j+1}$, we construct a binary word $w = \phi_d(m)$, having exactly d ones, as follows:

- Let $m_1 = m - 1$ and find p_1 such that $\binom{p_1}{d} \leq m_1 < \binom{p_1+1}{d}$;
- for every $2 \leq j \leq d$, let $m_j = m_{j-1} - \binom{p_{j-1}}{d-j+2}$ and find p_j such that
$$\binom{p_j}{d-j+1} \leq m_j < \binom{p_j+1}{d-j+1};$$
- starting from the right, make the binary word w of length n having a 1 at each position $p_j + 1$ for $j = 1, \ldots, d$, and adding as many zeros to the left as necessary.

For example, for $n = 8$, $d = 3$, and $m = 23$, we get $p_1 = 6$, $p_2 = 3$, and $p_3 = 0$, leading to the binary word $\phi_3(23) = 01001001$.

Note that if $p_1 + 1, \ldots, p_d + 1$ are the 1-positions associated with m and $q_1 + 1, \ldots, q_d + 1$ are the positions associated with k, then $m < k$ implies $\sum_{j=1}^{d} 2^{p_j} < \sum_{j=1}^{d} 2^{q_j}$. In other words, the corresponding words $\phi_d(m)$ and $\phi_d(k)$ are distinct, hence ϕ_d is one-to-one.

Given a binary word w of length n, having d ones, the inverse $m = \phi_d^{-1}(w)$ can be found as follows:

- Label all of the characters of w from right to left as $0, 1, 2, \ldots, n-1$;
- label the 1's in w from right to left as $1, 2, \ldots, d$ and record their positions as $p_1 + 1, \ldots, p_d + 1$ from left to right;
- define $m = 1 + \sum_{j=1}^{d} \binom{p_j}{d-j+1}$, with the convention that $\binom{a}{b} = 0$ if $a < b$.

For example, for the binary word $w = 01001001$, we get

$$w = \overset{7}{\mathbf{0}}\ \overset{\binom{6}{3}}{\boxed{\mathbf{1}}}\ \overset{5}{\mathbf{0}}\ \overset{4}{\mathbf{0}}\ \overset{\binom{3}{2}}{\boxed{\mathbf{1}}}\ \overset{2}{\mathbf{0}}\ \overset{1}{\mathbf{0}}\ \overset{\binom{0}{1}}{\boxed{\mathbf{1}}}$$

and therefore $\phi_3^{-1}(01001001) = 1 + \binom{6}{3} + \binom{3}{2} + \binom{0}{1} = 20 + 3 + 0 = 23$. □

With the help of ϕ_d and ϕ_d^{-1} we now proceed to prove the above proposition.

Proof of Proposition 7. Let $\alpha = (n_1^{c_1}, \ldots, n_k^{c_k})$ be an element of $A_k(\nu)$, where each $n_i^{c_i}$ is a part of size n_i with color $1 \leq c_i \leq \binom{n_i+d-1}{d}$. For every part $n_i^{c_i}$ let $w_i = \phi_d(c_i)$ be the binary word of length $n_i + d - 1$ obtained through the algorithm from Lemma 8. We then concatenate the k binary words associated with each part of the composition α, adding an extra 1 between consecutive parts, to create a binary string $\beta = T(\alpha)$ of length

$$(k-1) + \sum_{i=1}^{k}(n_i + d - 1) = k - 1 + \nu + (d-1)k = \nu + dk - 1$$

with exactly $(k-1) + dk = (d+1)k - 1$ ones. In other words, β is an element of $B_k(\nu)$.

For example, for $d = 2$ each part n takes on $d_n(2) = \binom{n+1}{2}$ colors, so for $n = 1, 2, 3$ the above map $T(\alpha) = \beta$ gives the following:

$d_n(2)$-color comp.	binary word
(1_1)	11

$d_n(2)$-color comp.	binary word
(2_1)	011
(2_2)	101
(2_3)	110
$(1_1, 1_1)$	11111

$d_n(2)$-color comp.	binary word
(3_1)	0011
(3_2)	0101
(3_3)	0110
(3_4)	1001
(3_5)	1010
(3_6)	1100
$(2_1, 1_1)$	011111
$(2_2, 1_1)$	101111
$(2_3, 1_1)$	110111
$(1_1, 2_1)$	111011
$(1_1, 2_2)$	111101
$(1_1, 2_3)$	111110
$(1_1, 1_1, 1_1)$	11111111

The above map T is reversible. Given a binary string β in $B_k(\nu)$, we create a composition in $A_k(\nu)$ by means of the following inductive algorithm:

- Write β as $w_1 1 \beta'$, where w_1 is a binary string with exactly d ones;
- let $c_1 = \phi_d^{-1}(w_1)$ and let $n_1^{c_1}$ be the part of size $n_1 = \text{length}(w_1)$ with color c_1;
- remove the one after w_1 and repeat the algorithm with β' until it has only d ones.

Since every such $\beta \in B_k(\nu)$ has exactly $(d+1)k - 1$ ones, the above algorithm will create k parts with $n_1 + \cdots + n_k = \nu$, leading to a composition $\alpha = (n_1^{c_1}, \ldots, n_k^{c_k})$ in $A_k(\nu)$ such that $T(\alpha) = \beta$. □

Let $\mathscr{A}_{p(d)}(\nu) = \bigcup_{k=1}^{\nu} A_k(\nu)$ be the set of $p(d)$-color compositions of ν.

Map $\mathscr{A}_{p(d)}(\nu) \to \mathscr{C}_{1,d+1}\big((d+1)\nu - 1\big)$. For $\alpha = (n_1^{c_1}, \ldots, n_k^{c_k})$ in $\mathscr{A}_{p(d)}(\nu)$, let $\beta = T(\alpha)$ be the binary string of length $\nu + dk - 1$ from Proposition 7, having exactly $(d+1)k - 1$ ones and $\nu - k$ zeros. If we treat every character 1 in β as a separate part and replace every 0 by $d+1$, we get a unique composition of $(d+1)k - 1 + (\nu - k)(d+1) = (d+1)\nu - 1$, having only parts of size 1 and $d+1$. This map is clearly a bijection.

For example, for $d = 2$ and $\nu = 3$, we get

$d_n(2)$-color comp.	binary word	comp. in $\mathscr{C}_{1,3}(8)$
(3_1)	0011	(3,3,1,1)
(3_2)	0101	(3,1,3,1)
(3_3)	0110	(3,1,1,3)
(3_4)	1001	(1,3,3,1)
(3_5)	1010	(1,3,1,3)
(3_6)	1100	(3,3,1,1)
$(2_1, 1_1)$	011111	(3,1,1,1,1,1)
$(2_2, 1_1)$	101111	(1,3,1,1,1,1)
$(2_3, 1_1)$	110111	(1,1,3,1,1,1)
$(1_1, 2_1)$	111011	(1,1,1,3,1,1)
$(1_1, 2_2)$	111101	(1,1,1,1,3,1)
$(1_1, 2_3)$	111110	(1,1,1,1,1,3)
$(1_1, 1_1, 1_1)$	11111111	(1,1,1,1,1,1,1,1)

Map $\mathscr{A}_{p(d)}(\nu) \to \mathscr{C}_{\equiv 1(d+1)}\big((d+1)\nu\big)$. For $\alpha = (n_1^{c_1}, \ldots, n_k^{c_k})$ in $\mathscr{A}_{p(d)}(\nu)$, let $\beta = T(\alpha)$ be the binary string of length $\nu + dk - 1$ from Proposition 7, having exactly $(d+1)k - 1$ ones and $\nu - k$ zeros. Using the 1's in β as separators, we now construct a composition as follows: To the left and right of every 1 in the binary string β, replace a string of j zeros with a string of $(d+1)j+1$ zeros, which then represents a part of size $(d+1)j+1$. Since there are $(d+1)k - 1$ separators, the constructed composition has $(d+1)k$ parts and the new total number of zeros is $(d+1)(\nu - k) + (d+1)k = (d+1)\nu$.

In conclusion, the above (clearly reversible) construction gives a composition of $(d+1)\nu$ having only parts of size congruent to 1 modulo $d+1$.

For example, for $d = 2$ and $\nu = 3$, we get

$d_n(2)$-color comp.	binary word	zeros as parts	comp. in $\mathscr{C}_{\equiv 1(3)}(9)$
(3_1)	0011	00000001010	(7,1,1)
(3_2)	0101	00001000010	(4,4,1)
(3_3)	0110	00001010000	(4,1,4)
(3_4)	1001	01000000010	(1,7,1)
(3_5)	1010	01000010000	(1,4,4)
(3_6)	1100	01010000000	(1,1,7)
$(2_1, 1_1)$	011111	00001010101010	(4,1,1,1,1,1)
$(2_2, 1_1)$	101111	01000010101010	(1,4,1,1,1,1)
$(2_3, 1_1)$	110111	01010000101010	(1,1,4,1,1,1)
$(1_1, 2_1)$	111011	01010100001010	(1,1,1,4,1,1)
$(1_1, 2_2)$	111101	01010101000010	(1,1,1,1,4,1)
$(1_1, 2_3)$	111110	01010101010000	(1,1,1,1,1,4)
$(1_1, 1_1, 1_1)$	11111111	01010101010101010	(1,1,1,1,1,1,1,1,1)

Map $\mathscr{A}_{p(d)}(\nu) \to \mathscr{C}_{\geq d+1}\big((d+1)\nu + d\big)$. For $\alpha = (n_1^{c_1}, \ldots, n_k^{c_k})$ in $\mathscr{A}_{p(d)}(\nu)$, let $\beta = T(\alpha)$ be the binary string of length $\nu + dk - 1$ from Proposition 7, having exactly $(d+1)k - 1$ ones and $\nu - k$ zeros. Using now the 0's in β as separators, we construct a composition as

follows: To the left and right of every 0 in the binary string β, replace a string of j ones with a string of $j+d+1$ ones, which then represents a part of size $j+d+1$. Since there are $\nu-k$ separators, the constructed composition has $\nu-k+1$ parts and the new total number of ones is $(d+1)k-1+(d+1)(\nu-k+1)=(d+1)\nu+d$.

Thus the above reversible construction gives a composition of $(d+1)\nu+d$ having no parts of size less than $d+1$.

For example, for $d=2$ and $\nu=3$, we get

$d_n(2)$-color comp.	binary word	ones as parts	comp. in $\mathscr{C}_{\geq 3}(11)$
(3_1)	0011	1110111011111	(3,3,5)
(3_2)	0101	1110111101111	(3,4,4)
(3_3)	0110	1110111110111	(3,5,3)
(3_4)	1001	1111011101111	(4,3,4)
(3_5)	1010	1111011110111	(4,4,3)
(3_6)	1100	1111101110111	(5,3,3)
$(2_1,1_1)$	011111	111011111111	(3,8)
$(2_2,1_1)$	101111	111101111111	(4,7)
$(2_3,1_1)$	110111	111110111111	(5,6)
$(1_1,2_1)$	111011	111111011111	(6,5)
$(1_1,2_2)$	111101	111111101111	(7,4)
$(1_1,2_3)$	111110	111111110111	(8,3)
$(1_1,1_1,1_1)$	11111111	11111111111	(11)

References

[1] M. Abrate, S. Barbero, U. Cerruti, N. Murru, Colored compositions, invert operator and elegant compositions with the "black tie", *Discrete Math.* **335** (2014), 1–7.

[2] M. Bernstein and N. J. A. Sloane, Some canonical sequences of integers, *Linear Algebra Appl.* **226/228** (1995), 57–72.

[3] L. Comtet, *Advanced Combinatorics: The Art of Finite and Infinite Expansions*, D. Reidel Publishing Co., Dordrecht, 1974.

[4] C. R. Dedrickson, "Compositions, Bijections, and Enumerations," (2012), Electronic Theses & Dissertations, Paper 17.

[5] S. Eger, Identities for partial Bell polynomials derived from identities for weighted integer compositions, *Aequat. Math.*, March 2015.

[6] S. Heubach and T. Mansour, *Combinatorics of Compositions and Words*, Discrete Mathematics and its Applications (Boca Raton), CRC Press, Boca Raton, FL, 2010.

[7] V. E. Hoggatt, D. A. Lind, Fibonacci and binomial properties of weighted compositions, *J. Combin. Theory* **4** (1968), 121–124.

[8] L. Moser and E. L. Whitney, Weighted compositions, *Canad. Math. Bull.* **4** (1961), 39–43.

[9] C. Shapcott, New bijections from n-color compositions, *J. Comb.* **4** (2013), no. 3, 373–385.

Department of Mathematics, Nazareth College, 4245 East Ave., Rochester, NY 14618

Penn State Altoona, 3000 Ivyside Park, Altoona, PA 16601